



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

the excavated soil afterward returned. They were square, 3 feet each way at top, 4 feet deep, and open at bottom. Sterilizing the soil was done by a Company railway engine, the soil in each pot being steamed two hours, at 110° C. Fourteen pots were thus sterilized and the remaining six were neither sterilized nor inoculated. All the pots were planted with one "bit" of a banana tuber having two "eyes" and in seven of those sterilized these bits were inoculated with a laboratory culture of *Fusarium Cubense*, while the other seven were not inoculated. The inoculum was a combination of two kinds of culture—mixed at time of application—one in corn meal decoction and the other in Uschinski's solution, both originating from rice tube cultures, three weeks previously. It was applied in quantity of about one liter to each pot. Already in February, 1922, one of the inoculated plants showed signs of disease. By April 1 every one of the inoculated plants was diseased and most of them were even then bad cases. Up to July 12, when the writer left Panama, no one of the plants in the inoculated pots, either sterilized or unsterilized, gave any indication of disease. Besides the added interest that may be attached to this experiment, with a different variety of banana in a different locality, the writer is glad, and others will be, that it confirms the painstaking work of Dr. Brandes. It should be added that Dr. J. R. Johnston, director of tropical research for the United Fruit Company proposed the experiment, in which the writer was also supported and encouraged by the manager, H. S. Blair, and by H. L. Peck, E. C. Adams, and other men of the Company.

MARK ALFRED CARLETON

CUYAMEL, HONDURAS

FRESH WATER COELENTERATA IN KENTUCKY

In 1916¹ the writer called attention to an extraordinary appearance of the rare fresh-water medusa, *Craspedacusta sowerbyi*, in a Kentucky creek, when it was present literally by the thousands, and expressed his intention to make a search for the hydroid stage of the

jellyfish, since the indications were that where so many of the sexual form occurred there should be a good prospect of finding the asexual stage. With this in view visits have been made from time to time to the place where the medusæ were found. In 1917, a year after they were first observed, they were present again in as great numbers as in the preceding year²; still no trace of a hydroid condition could be found. Each year thereafter the place was searched, but they have not been seen since 1917. It may be of interest to record, however, that this year (September 26) when examining the locality, some fine growths of the hydroid *Cordylophora*³ were obtained in the part of the stream where *Craspedacusta* was discovered. There can be no doubt about this material representing *Cordylophora lacustris*, and this interesting fresh-water hydroid is thus given a new locality in which it is associated in a way with *Craspedacusta*. The growths are attached to living willow twigs that by the drooping of branches have become immersed in the water.

HARRISON GARMAN

UNIVERSITY OF KENTUCKY
LEXINGTON, KY.

AN ANNOUNCEMENT IN SCIENCE

TO THE EDITOR OF SCIENCE: I am glad to be able to inform you that in response to my appeal on behalf of Professor W. Boldyreff, published in SCIENCE of October 27 under the heading, "An Opportunity," I received promptly a number of requests for special information concerning Dr. Boldyreff's qualifications, and that as a result of these inquiries he has now received an appointment for a year in a dignified position which will give him excellent opportunity for research work and an income to keep himself and family alive.

VERNON KELLOGG

NATIONAL RESEARCH COUNCIL,
WASHINGTON, D. C.,
NOVEMBER 28, 1922

² Professor Hargitt in commenting on the matter (SCIENCE, L. 1919, page 413) was wrong in inferring that they disappeared at once.

³ A medusa stage is believed not to be produced by this hydroid.

¹ SCIENCE, Vol. XLIV, page 858.